

Testicular yolk sac tumor in an eight-month old child: A case report

Khan IU¹, Jose J², Fawazy T¹, Hadi WA², Sharma PK¹

ABSTRACT

Germ cell tumors are rare in children. This group of tumors in children is distinct from that in adults in terms of their behavior, histological nature, metastasis and the treatment needed. 85% of the yolk sac tumors in children present as clinical stage I disease as compared to 35% in adults. AFP can be used as a reliable tumor marker as the levels are increased in more than 90% of yolk sac tumors. In children inguinal orchidectomy followed by stringent surveillance for two years is the preferred treatment option. Combination chemotherapy may be reserved for metastatic disease or recurrence. Retroperitoneal Lymph Node Dissection (RPLND) is preferable in adults. In this case report we describe the clinical presentation, imaging findings, diagnosis and up-to-date management of yolk sac tumor in an eight-month old child.

Key words: germ cell tumor, yolk sac tumor, Alpha fetoprotein, chemotherapy, RPLND

Citation

Khan IU, Jose J, Fawazy T, Hadi WA, Sharma PK. Testicular yolk sac tumor in an eight-month old child: A case report. Gulf Medical Journal. 2012;1(1):37-40.

INTRODUCTION

Testicular tumors are rare in children, but highly treatable and usually curable. In children testicular tumors account for approximately 1 to 2% of all tumors, with an incidence of 0.05-2 per 100, 000 children. Among testicular tumors germ cell tumors have a bimodal age distribution. One peak occurs in the first two years of life and the second in young adults 15-35 years of age. Pediatric testicular tumors are dramatically different from adult neoplasms. Germ cell tumors account for 60% testicular tumors in children, but 95% of testicular tumors in adults. Adult germ cell tumors like seminoma and embryonal carcinoma are rare in children. Teratoma which are benign in children are often malignant in adults.

This is the first case of testicular cancer, yolk sac tumor, in an eight-month old child in the author's 27 years of experience in urology.

CASE REPORT

A mother brought an eight-month old child to the Urology OPD with a painless right testicular swelling, which was noticed one week previously. This was the first clinic that the child had attended. On examination it was 6x5cm in size, non-tender, irregular in shape and firm in consistency.

There was a family history of cancer. The uncle of child had died at the age of 18 years because of leukemia a few years previously.

Color Doppler ultrasound examination revealed an enlarged right testis measuring 4x3x2.5cm with estimated volume of 16.3 cc. It showed multiple hypoechoic cystic foci of different sizes and appeared hypervascular on color Doppler evaluation. Hypervascularity was also noticed within the right spermatic cord. A mild amount of free fluid was detected in the right scrotal compartment. As the normal size of the left testicle for this age is about 1.7x1.1x0.6 cm with an estimated volume of 0.56 cc., the clinical diagnosis made was enlarged right testicle, possibly neoplastic in nature.

MRI of the upper abdomen, pelvis and male perineum revealed a diffusely enlarged right testis measuring 4.6x2.8x4.4 cm, displaying

¹Department of Urology, Gulf Medical College Hospital & Research Center, ²Dept. of Pathology, Gulf Medical University, Ajman, UAE. Correspondence: Dr. Ihsan Ullah Khan, Department of Urology, Gulf Medical College Hospital & Research Center, Ajman, UAE. email: miukhan@hotmail.com

heterogenous high signal in T2 and fat suppressed images with central necrosis and appreciating heterogenous yet significant post contrast enhancement, a pattern that is indicative of a neoplastic process. The ipsilateral epididymis was diffusely enlarged with significant post contrast enhancement. Mildly thickened and enhancing ipsilateral spermatic cord. The right scrotal compartment was distended by the diffusely infiltrated testis with associated minimal hydrocele and diffuse thickening and enhancement of the surrounding skin. The left testis was displaced in an anterior and superior direction by the mass effect and noted in the left scrotal neck and measuring 16x8 mm in diameter. A few bilateral inguinal LN with no suspicious criteria were observed. Additionally, liver, spleen, gall bladder, pancreas, kidneys, major vessels and urinary bladder were found to be normal. No detectable masses or lymph nodes were seen in the abdomen. Both lung fields and C/P angles were clear on chest X-ray.

The values of tumor markers Alpha Fetoprotein (AFP) and beta HCG are shown in Table 1.

The patient underwent unilateral orchidectomy through inguinal approach. The right testis was removed together with all its coverings with maximum cord length up to the deep inguinal ring. Spermatic veins were observed to be dilated and tortuous. No fixity to skin was seen. The patient was discharged the next day and stitches removed

Table 1. Values of tumor markers

| Tumor Marker | Observed Value | Biological Reference Range |
|--------------|----------------|----------------------------|
| AFP | 29618 ng/ml | 0.2-9.0 ng/ml |
| Beta HCG | 0.500 miu/ml | Less than 5 miu/ml |

after eight days. The post-operative period was uneventful.

Gross examination of the specimen submitted for histopathology revealed a well circumscribed gray-white neoplasm 4x4x3 cm, with areas of necrosis and cystic change, almost totally replacing testicular parenchyma (Figure 1).

H&E stained sections of the tumor showed cuboidal tumor cells arranged mostly in a reticular-microcystic pattern. The cells exhibited pleomorphism, hyperchromatisia and frequent mitosis with occasional intracellular and extracellular eosinophilic hyaline globules. Glomeruloid papillary structures composed of



Figure 1. Cut section of the testicular tumor revealing a circumscribed solid gray-tan neoplasm with necrosis and cystic change

tumor cells covering a central fibrovascular core within a cystic space lined by a single layer of flat/cuboidal cells (Schiller-Duval Bodies) were seen in some areas (Figure 2). Solid and microcystic patterns were also observed. No seminomatous, choriocarcinomatous or teratomatous elements were identified. A few compressed seminiferous tubules were present at the periphery of the tumor. Sections from the epididymis, spermatic cord and rete testis were free of tumor.




Figure 2. Schiller-Duval body (arrow) characterized by tumor cells forming glomeruloid papillary structure is pathognomonic of yolk sac tumor. (H&E x20)

Immunohistochemical staining was negative for CD30 and focally positive for AFP. The findings were consistent with pure yolk sac tumor. The patient is under stringent follow-up. AFP analysis done 17 days after surgery had dropped significantly to 636 ng/ml from its 29618 ng/ml before surgery. He was referred to the oncologist for further management and follow up.

DISCUSSION

Germ cell tumors are most frequently found in the gonads. Only 2-5% of them arise in extragonadal regions such as mediastinum, retroperitoneum, pineal gland and sacral areas. Yolk sac tumor of penile shaft and urachus has also been reported².

Germ cell tumors are of two types, seminomas and non-seminomas. Seminomas are extremely rare in prepubertal children. Among non-seminomatous germ cell tumors the most common are teratomas and yolk sac tumors which account for about 62% and 26% of testes tumor in infants and toddlers respectively³.

Teratomas are described in some series as constituting 50% of prepubertal testicular tumors. However these are reported less because of their benign nature⁵. In tumor registries, yolk sac tumors are mentioned more commonly than teratomas¹.

Different risk factors have been described for testicular cancer in the early age group. They include pre-natal exposure to high estrogen levels during intrauterine life, gonadal dysgenesis, exposure of mother to chemical carcinogens and smoking, trauma, testicular atrophy due to torsion, and orchitis.

Family history of cancer is an important risk factor in the early age groups. In the case of this child the uncle had died of leukemia at around 18 years of age. However no other causal relationship has been established.

Three clinical stages for the determination of extension of the tumor have been described⁴. Stage I is where the tumor is confined to the testis. Invasion of epididymis tunica albuginea, spermatic cord or scrotum does not change tumor stage but increases the risk of nodal involvement and the risk of recurrence.

In stage II the tumor has retroperitoneal lymph node metastasis. Stage III is characterized by supraclavicular lymph nodes, visceral involvement or persistently elevated tumor marker values.

The vast majority, 85%, of yolk sac tumors in children present as clinical stage I disease compared to 35% in adults¹. AFP can be used as a reliable tumor marker as levels are increased in more than 90% of yolk sac tumors. Therefore patients can be safely managed with observation after orchidectomy. Chemotherapy is reserved for recurrence. In our case, the level of tumor marker AFP was 2,9618ng/ml before surgery. MRI revealed no lymph node involvement retroperitoneally. X-ray chest was also clear. Our

patient was in stage IC-T1 No Mo S3.S3 means AFP more than 10,000 ng/ml.

Tumor marker AFP dropped significantly to 636ng/ml on 17th post-operative day.

The nature of yolk sac tumor is different in children from that in post-pubertal adults. In young children yolk sac tumors are predominantly pure in histology and are not usually mixed with other histological types as may be encountered in adults. Histopathology and immunohistochemistry revealed that our patient had pure yolk sac tumor.

Yolk sac tumors in children metastasize through the hematogenous route in more than half of the cases in comparison to adults where it is only 4-6%⁵. This fact changes the treatment modality. It indicates that RPLND would not be adequate treatment in children and would result in complications like chylous ascities, post-operative bowel obstruction, wound infection and subsequently ejaculatory dysfunction.

The German Society of Pediatric Oncology reported that 15% of patients had recurrence with stage I yolk sac tumor after radical inguinal orchidectomy and observation alone. These patients were cured successfully by combination chemotherapy, with 100% survival. Even in the patients with metastatic disease that they treated, four out of five had remission with chemotherapy⁵.

Keeping the above in view we recommended observation after inguinal orchidectomy and stringent follow-up for two years for our patient. The latter includes tumor marker AFP, x-ray chest every two months and MRI every 3-4 months. If there is no evidence of recurrence after the second year, surveillance using all these methods can be less frequent since the risk is highest in the first two years.

CONCLUSION

Pre-pubertal testicular tumors are rare but result in great anxiety and burden on the family when they do occur. Testicular germ cell tumors in infants are distinct from those in adults. In recent years the prognosis of testicular tumors, even in metastatic disease, has changed dramatically with combination chemotherapy⁸: In stage I 99-100%; In stage II 95-98%; In stage III 45%-75%. Despite this good prognosis, the parents have many reservations regarding the treatment and its effects on the child's growth and particularly on future fertility.

References

1. Christopher SC. Prepubertal testicular and paratesticular tumors: Medscape reference.2011 web MD, LLC.
2. Samaila, Modupeola, Maitama HY, Abdullahi K, Mbibu H, Waziri GD. Yolk sac tumor of the penile shaft: A rare primary extra-gonadal presentation. *Afr J Paediatr Surg.* 2011;8:241-3.
3. Huang H, KO SF, Chuang JH, Jeng YM, Sung MT, Chen WJ. Pprimary yolk sac tumor of the urachus; *Arch Pathol Lab Med.* 2002;126:1106-9.
4. Yilmaz A, Bayraktar B, Sagiroglu J, Gucluer B. A case report: Giant seminoma in an undescended testis presenting as an abdominal wall mass *journal of surgical case reports-ISSN:2042-8812-2011.12:9.*
5. Ahmed HU, Arya M, Muneer A, Mushtaq I, Sebire NJ. Testicular and paratesticular tumors in the prepubertal population. *Lancet Oncol.* 2010;11:476-83.
6. Testicular cancer in children:2012 AUA Foundation:Jan.2011. Available from: URL: <http://www.urologyhealth.org/>
7. Kim T, Kim DY, Cho MJ, Kim SC, Seo JJ, Kim IK. Use of Kim T, Kim DY, Cho MJ, Kim SC, Seo JJ, Kim IK. Use of laparoscopic surgical resection for pediatric malignant solid tumors: a case series; *Surg Endosc.* 2011; 25:1484-8.
8. Presti JC Jr. Genital tumors. 17th ed. In: Tanagho EA, McAnich JW, editors. *Smith's general urology.* New York: McGraw Hill. 2008. p. 375-87.